

A method to illustrate the extensive and intensive properties of thermodynamic variables

Stephen R. Addison
Department of Physics and Astronomy
University of Central Arkansas
Conway, AR 72035

INTRODUCTION

Over the last ten years, unified courses in thermal physics, including both macroscopic thermodynamics and microscopic statistical mechanics, have become increasingly common, particularly at the undergraduate level. This reflects a change from traditional curricula, where entire courses are devoted to classical thermodynamics and statistical physics is briefly introduced in survey courses in modern physics with serious study delayed until graduate school. A thorough understanding of statistical and probabilistic is becoming increasingly important to chemists and physicists. This understanding is often developed by a serious study of statistical mechanics. Undergraduate thermal physics provides an ideal foundation for such study.

Undergraduate courses in thermal physics do not afford us the time to follow the languid, historical, phenomenological development of the laws of thermodynamics as exemplified in the classic text by Zemansky (1981). The study of thermodynamics can be abbreviated by using the axiomatic method. The most widely accepted axioms are those proposed by Callen (1960). A thorough discussion of these axioms is provided in Callen's *Thermodynamics* and in the more recent book by Tien and Lienhard (1985). In the axiomatic approach, the extensive or intensive nature of thermodynamic variables is emphasized and the relationship to the theory of homogeneous functions is presented. However, this relationship is not often exploited, and an opportunity to provide a foundation for the later study of critical point phenomena is lost.

In the following sections the relationship between homogeneous function theory and thermodynamics is developed, and, as an illustration, a method for calculating the thermodynamic properties of N moles of a material if an equation is known for a fixed amount of that material. The author first became familiar with some of these methods in a course based on Cullen's text, and began to use them some years later after having developed a course in thermal physics. An exhaustive literature search has failed to find a description of the methods described herein.

HOMOGENEOUS FUNCTIONS

As the properties of homogeneous functions are not well known, their properties are reviewed here. As is well known, a polynomial of the form

$$A_0 + A_1x + A_2x^2 + \dots + A_nx^n \quad (1)$$

is of degree n if A_n is not equal to zero. Thus, the degree of a polynomial is equal to the largest exponent in the polynomial. A polynomial in more than one variable is said to be *homogeneous* if all of its terms are of the same degree. For example, the polynomial in variables x and y

$$f(x, y) = x^2 + 2xy + y^2 \quad (2)$$

is homogeneous of degree 2. This much is familiar.

The property of homogeneity can be extended to more general functions. We shall consider general functions that are homogeneous in terms of functional equations. The functional equations appropriate to the study of homogeneous functions were developed by Euler (1755, 1768, 1770). Aczel (1966, 1969), Davis (1962), Stanley (1970), and Widder (1961) provide modern introductions with varying degrees of sophistication to the subject of functional equations.

In general, a function $f(x)$ is a homogeneous function if for all values of the parameter λ ,

$$f(\lambda x) = g(\lambda)f(x) \quad (3)$$

The function g is usually called the scaling function in thermodynamics. The function g is not an arbitrary function, it is given by

$$g(\lambda) = \lambda^n, \quad (4)$$

see for example Stanley (1971).

Thus a homogeneous function $f(x)$ is one that satisfies

$$f(x\lambda) = \lambda^n f(x) \quad (4)$$

This definition can be extended to any finite number of variables. In elementary thermodynamics we are usually interested in functions that are homogeneous of degree zero or one. It is possible for multidimensional functions to be homogeneous in different degrees for the different variables. This is common occurrence in thermodynamics. For a function in the variables x , y , and z ; the function

$$f(\lambda x, \lambda y, z) = \lambda^n f(x, y, z) \quad (5)$$

then we say that this function is homogeneous of degree n in x and y and that it is not homogeneous in z .

A generalization, described by Stanley (1971), is that of a *generalized homogeneous function*. Generalized homogeneous functions are those that satisfy

$$f(\lambda^a x, \lambda^b y) = \lambda f(x, y). \quad (6)$$

It is this formulation that is widely used in the analysis of critical point phenomena and phase transitions using the static scaling hypothesis.

For clarity, the homogeneities of degree zero and one are treated separately. The reader will readily observe that a single theorem,

true for any n , could have been presented. The method of proof is a straightforward extension of the proofs presented below. The author has presented both methods in thermal physics courses, and has found the separate development presented below is preferred by most students. The unified theorem is assigned as a homework problem.

In thermodynamics, variables are classified as either *extensive* or *intensive*. Extensive variables are those, which vary linearly with the size of the system. Internal energy, E , is an example of an extensive variable. Extensive variables exhibit the property of being additive over a set of subsystems. As example: if a system is composed two subsystems, one with energy E_1 , the second with energy E_2 , then the total system energy is $E = E_1 + E_2$. Other examples of extensive variables in thermodynamics are: volume, V , mole number, N , entropy, S , enthalpy, H , Helmholtz free energy, F , and the Gibbs free energy, G .

Other thermodynamic variables are independent of the size of the system, these variables are referred to as *intensive* variables. Examples of intensive variables encountered in thermodynamics are temperature, T , pressure, p , and the *chemical potential* of the i^{th} component of the system, μ_i .

THE RELATIONSHIP BETWEEN HOMOGENEOUS FUNCTIONS AND EXTENSIVE AND INTENSIVE VARIABLES

Entropy, which is usually an extensive variable (there are exceptions – see Hill (1962), Landsberg (1978), or Robertson (1993)) in thermodynamics, can be expressed as a function of three other extensive variables: internal energy, volume, and number of moles. Thus, we may write entropy as $S = S(E, V, N)$. An equation of this form contains complete thermodynamic information, it is called the *fundamental relation* in thermodynamics. Since extensive variables depend on the size of the system, they can be represented by

homogeneous first order degree functions. Given the fundamental relation for entropy, we may write

$$S(\lambda E, \lambda V, \lambda N) = \lambda S(E, V, N). \quad (7)$$

The scaling function λ is arbitrary, and following Callen (1960) we choose $\lambda = 1/N$. The entropy equation can now be rewritten as

$$S\left(\frac{E}{N}, \frac{V}{N}, 1\right) = \frac{1}{N} S(E, V, N). \quad (8)$$

Next we introduce the internal energy per mole as $e = E/N$, and the volume per mole $v = V/N$, then

$$S(e, v, 1) = s(e, v) = \frac{1}{N} S(E, V, N) \quad (9)$$

or

$$Ns(e, v) = S(E, V, N) \quad (10)$$

where $s(e, v)$ is the entropy per mole. This tells us that the entropy of N moles of a substance is N times the entropy per mole of that substance. The result is well known, the method used to derive it can be generalized to provide a result of more general utility.

The intensive functions of thermodynamics are homogeneous functions of order zero of the extensive variables. Callen (1960) has shown that the intensive parameters of a thermodynamic systems can be written as a function of the extensive variables; that is $T = T(S, V, N)$, $P = P(S, V, N)$, and $\mu = \mu(S, V, N)$. These expressions are called *equations of state*. Each of these functions is homogeneous of degree zero, this follows as each of these can be expressed as a first partial derivative of the energy or entropy of the system. The details can be found in Chandler (1987). For example, the temperature function $T = T(S, V, N)$ obeys

$$T(\lambda S, \lambda V, \lambda N) = T(S, V, N). \quad (11)$$

This means that in a composite system in thermal equilibrium, the temperature in any sub-subsystem is equal to the temperature of the system.

Two Theorems About Intensive and Extensive Functions and their Application

We now establish several results on intensive and extensive functions and then discuss some thermodynamic applications of them.

THEOREM 1 (Extensive Functions): Consider a function $f(x, y, z)$, homogeneous of degree one in the variables y and z . Let one of the homogeneous variables be a constant, $z = z_0$, then we can write $f(x, y, z_0) = g(x, y)$. Then

$$f(x, y, z) = \frac{z}{z_0} g(x, y \frac{z}{z_0}). \quad (12)$$

PROOF: Write $f(x, y, z)$ as

$$f(x, y, z) = \frac{z}{z_0} \frac{z_0}{z} f(x, y, z). \quad (13)$$

Then, by choosing $\lambda = \frac{z}{z_0}$, we can rewrite (13), using the fact it is homogeneous in the variables y and z , so we have

$$f(x, y, z) = \frac{z}{z_0} f(x, y \frac{z_0}{z}, z \frac{z_0}{z}) = \frac{z}{z_0} f(x, y \frac{z_0}{z}, z_0) \quad (14)$$

Now, since z_0 is a constant, we can write (14) as

$$f\left(x, y \frac{z}{z_0}, z_0\right) = g\left(x, y \frac{z_0}{z}\right), \quad (15)$$

thus, we have

$$f(x, y, z) = \frac{z}{z_0} g\left(x, y \frac{z}{z_0}\right) \quad (16)$$

which completes the proof.

THEOREM 2 (Intensive Functions) : Consider a function $f(x, y, z)$, that is homogeneous degree zero in the variables y and z . Let one of the homogeneous variables be a constant, $z = z_0$, then we can write $f(x, y, z_0) = h(x, y)$. Then

$$f(x, y, z) = h\left(x, y \frac{z}{z_0}\right). \quad (17)$$

PROOF: Write $f(x, y, z)$ as

$$f\left(x, y \frac{z_0}{z}, z_0\right) = f(x, y, z) \quad (18)$$

since it is zero order homogeneous in y and z , then

$$h\left(x, y \frac{z_0}{z}\right) = f(x, y, z) \quad (19)$$

which is the theorem.

We will now consider some illustrations of these theorems. Given that the internal energy of a system for three moles of a substance is

$E = pV^2$, what is the internal energy for N moles? We want $E(p, V, N)$, using Theorem 1 (noting that p is intensive, V and N extensive)

$$E(p, V, N) = \frac{N}{N_0} g(p, V \frac{N_0}{N}), \quad (20)$$

where $g(p, v) = pV^2$, and $N_0 = 3$. Thus,

$$E(p, V, N) = \frac{N}{3} (P(\frac{3V}{N})^2) = 3 \frac{pV^2}{N}. \quad (21)$$

As a second illustration of Theorem 1, consider the entropy of one mole of some substance that has the equation of state

$$s = R \ln(E^{3/2}V) + C, \quad (22)$$

where C is a constant and R is the ideal gas constant. To determine the entropy for N moles of this material, we use from Theorem 1 to write (22) as

$$S = \frac{N}{N_0} \left\{ R \ln \left(\left(\frac{EN_0}{N} \right)^{3/2} \frac{VN_0}{N} \right) + C \right\}, \quad (23)$$

but $N_0 = 1$, so

$$S = NR \ln(E^{3/2}VN^{-5/2}) + NC. \quad (24)$$

Many similar results could be proved, but these are sufficient to illustrate Theorem 1.

We now illustrate Theorem 2. Again, let the number of moles, N play the role of the coordinate z so that $z_0 = N_0$. The van der Waals equation of state for a single mole of gas is

$$p = \frac{RT}{V-b} - \frac{a}{V^2}, \quad (25)$$

where a and b are phenomenological constants. We use Theorem 2 to find the equation of state for N moles. Applying the theorem, we write

$$p(T, V, N) = p\left(T, V \frac{N_0}{N}\right), \quad (26)$$

so the pressure for N moles is

$$p = \frac{RT}{\frac{V}{N} - b} - \frac{a}{\left(\frac{V}{N}\right)^2} = \frac{NRT}{V - Nb} - \frac{N^2 a}{V^2}, \quad (27)$$

which is, of course, the van der Waals equation for N moles.

Discussion and Conclusions

This paper has demonstrated the usefulness of functional equations in thermodynamics. One of us has used the theorems in thermal physics courses taught over the years. A study of these theorems circumvents the usual tendency of students to scale thermodynamic results incorrectly. The theorems presented here provide a deeper understanding of the relationship between the fundamental relation of thermodynamics and the equations of state.

The methods of functional equations are extremely useful in the analysis of critical point phenomena, and while not well known, they can be and are being applied to other sub-fields of physics. Some examples are the application to relativity Lunn (1919), to communication theory by Shannon (1948), and to information theory by Jaynes (1957). These applications to a number of different fields

suggest that physicists should become more familiar with the techniques of applying functional equations. The theorems derived in this paper are useful for calculating general results from measurements made on fixed amounts of materials. The methods developed in this paper allow the student to develop an understanding of the mathematical techniques used in the application of homogeneous functions; this allows these students to concentrate on the physics of critical point phenomena when they are first met, thus affording a deeper understanding.

REFERENCES

1. J. Aczel, *Lectures on Functional Equations and Their Applications*, Academic Press, New York, 1966.
2. Aczel, *On Applications and Theory of Functional Equations*, Academic Press, New York, 1969.
3. Herbert B. Callen, *Thermodynamics*, Wiley, New York, 1960.
4. David Chandler, *Introduction to Modern Statistical Mechanics*, Oxford University Press, New York, 1987.
5. Harold T. Davis, *Introduction to Non-Linear Differential and Integral Equations*, Dover, New York, 1962.
6. Leopold Euler, *Institutiones Calculi Differentialis*, Berlin, 1775.
7. Leopold Euler, *Institutiones Calculi Integralis*, Petropol., 1768.
8. Leopold Euler, *Institutiones Calculi Differentialis III*, Petropol. 1770.
9. Terrel L Hill, Thermodynamics of small Systems, J. Chem. Phys. **36**, 3182-3197, (1962).
10. E. T. Jaynes, Information Theory and Statistical Mechanics I, Phys. Rev. **106**, 620-630, 1957.
11. Peter T. Landsberg, *Thermodynamics and Statistical Mechanics*, Oxford (1978).
12. Robertson, Harry S., *Statistical Thermophysics*, Prentice Hall (1993).
13. A. C. Lunn, Some Functional Equations in the History of Relativity, Bull. Am. Math. Soc. **21**, 26-24, 1919.

14. Claude E. Shannon, A Mathematical Theory of Communication I, Bell. System Tech. J. **27**, 379-423, 1948.
15. Claude E. Shannon, A Mathematical Theory of Communication II, Bell. System Tech. J. **27**, 623-655, 1948.
16. H. E. Stanley, *Introduction to Phase Transitions and Critical Phenomena*, Oxford University Press, New York, 1971.
17. C. L. Tien and J. H. Lienhart, *Statistical Thermodynamics*, Revised Printing, Hemisphere Publishing Corporation, New York, 1985.
18. D. V. Widder, *Advanced Calculus*, 2nd. Edition, Prentice-Hall, Englewood Cliffs, New Jersey, 1961.
19. M. W. Zemansky and R. H. Dittman, *Thermodynamics*, 6th Edition, McGraw-Hill, New York, 1981.